

A NOTE ON PRINCIPAL PARTS ON PROJECTIVE SPACE AND LINEAR REPRESENTATIONS

HELGE MAAKESTAD

ABSTRACT. Let H be a closed subgroup of a linear algebraic group G defined over a field of characteristic zero. There is an equivalence of categories between the category of linear finite-dimensional representations of H , and the category of finite rank G -homogeneous vector bundles on G/H . In this paper we will study this correspondence for the sheaves of principal parts on projective space, and we describe the representation corresponding to the principal parts of a line bundle on projective space.

1. INTRODUCTION

In this note we will study the vector bundles of principal parts $\mathcal{P}^k(\mathcal{O}(n))$ of a line bundle on projective space over a field F of characteristic zero from a representation theoretic point of view. We consider projective N -space as a quotient $SL(V)/P$, where V is an $N + 1$ -dimensional vectorspace over F , and P is the subgroup of $SL(V)$ stabilizing a line L in V . There is an equivalence of categories between the category of finite rank $SL(V)$ -homogeneous vector bundles on $SL(V)/P = \mathbf{P}(V^*)$ and the category of linear finite-dimensional representations of P . The principal parts $\mathcal{P}^k(\mathcal{O}(n))$ are $SL(V)$ -homogeneous vector bundles on $\mathbf{P}(V^*)$, and the novelty of this note is that we describe the P -representation corresponding to the principal parts. The main result is Theorem 2.4. The Theorem says the following: Let L^* be the dual of the P -module L . Then for all $1 \leq k < n$, the P -representation corresponding to $\mathcal{P}^k(\mathcal{O}(n))$ is $S^{n-k}(L^*) \otimes S^k(V^*)$. As a corollary, we obtain the splitting-type of $\mathcal{P}^k(\mathcal{O}(n))$ on $\mathbf{P}(V^*)$ for all $1 \leq k < n$, and recover results obtained in [5],[6], [7] and [8].

2. PRINCIPAL PARTS ON PROJECTIVE SPACE

In this section we give the representation corresponding to $\mathcal{P}^k(\mathcal{O}(n))$ on $\mathbf{P}(V^*)$ for all $1 \leq k < n$, where V is an F -vectorspace of dimension $N + 1$ and F is a field of characteristic 0. A variety is an integral scheme of finite type over F . We will consider closed points when we talk about points of a scheme. Let V be a finite dimensional vectorspace over F . We let $GL(V)$ denote the group of all invertible linear transformations of V . It is an algebraic group in the sense of [2], chapt. 1. A *linear algebraic group* is a closed subgroup of $GL(V)$. Let $SL(V)$ be the linear

Date: 2002.

Key words and phrases. Homogeneous spaces, homogeneous vectorbundles, principal parts, linear representations, splitting-type.

Partially supported by the Emmy Noether Research Institute for Mathematics, the Minerva Foundation of Germany, the Excellency Center "Group Theoretic Methods in the study of Algebraic Varieties" of the Israel Science foundation and the EAGER foundation (EU network, HPRN-CT-2000-00099).

algebraic group of linear transformations of V with determinant 1. Let L in V be a line, and P the closed subgroup of $SL(V)$ stabilizing L . Then the quotient $SL(V)/P$ (which exists by [2], Theorem 6.8) is isomorphic to $\mathbf{P}(V^*)$, the projective space of lines in V (see [1], Section 4.2. This works over any field, not only the complex numbers). There exists a natural left $SL(V)$ -action on $\mathbf{P}(V^*)$, making it into a *homogeneous space* for $SL(V)$. And by [1], Chapt. 4, there exists an equivalence of categories between the category of finite rank homogeneous vector bundles on $SL(V)/P$ and the category of linear finite-dimensional representations of P , and under this correspondence the dimension of the representation gives the rank of the corresponding vector bundle. Hence any character of P gives a homogeneous line bundle on $SL(V)/P$. The line L corresponds to a character of P , and the bundle corresponding to the dual line L^* , is the line bundle $\mathcal{O}(1)$ on $\mathbf{P}(V^*)$ (see [1], Section 4.2). It is also a standard fact that any linear finite-dimensional representation ρ of P lifting to a representation $\tilde{\rho}$ of $SL(V)$ corresponds to a trivial abstract vector bundle on $\mathbf{P}(V^*)$. There exists on any scheme an equivalence of categories between the category of locally free finite rank sheaves and the category of finite rank vectorbundles, hence we will use these two notions interchangeably.

Pick a basis $e_0, \dots, e_N$ for V . Let $x_0, \dots, x_N$ be the dual basis, and let L be the line spanned by e_0 . Having chosen a basis for V it follows that $SL(V)$ may be identified with the group of square rank $N+1$ matrices with determinant equal to 1. The group P may be identified with the subgroup of $SL(V)$ consisting of matrices g of the form

$$g = \begin{pmatrix} a & * & \cdots & * \\ 0 & a_{11} & \cdots & a_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & a_{n1} & \cdots & a_{nn} \end{pmatrix}.$$

The one dimensional representation $\chi : P \rightarrow GL(S^n(L^*))$ corresponding to the line bundle $\mathcal{O}(n)$ is given by $\chi(g) = a^{-n}$.

Let X be a smooth variety of dimension d and consider the diagonal Δ in $X \times X$. Let $\mathcal{I}$ be the sheaf of ideals of $\mathcal{O}_{X \times X}$ defining the diagonal Δ , and put $\mathcal{O}_{\Delta^k}$ to be $\mathcal{O}_{X \times X} / \mathcal{I}^{k+1}$.

Definition 2.1. Let p, q be the projection maps from $X \times X$ to X , and let $\mathcal{E}$ be an $\mathcal{O}_X$ -module. Define $\mathcal{P}^k(\mathcal{E}) = p_*(\mathcal{O}_{\Delta^k} \otimes q^*\mathcal{E})$ to be the k 'th order principal parts of the module $\mathcal{E}$. We put $\mathcal{P}^k(\mathcal{O}_X) = \mathcal{P}^k$.

Note that by [5], if the rank of $\mathcal{E}$ is e , $\mathcal{P}^k(\mathcal{E})$ is a vector bundle of rank $e \binom{d+k}{d}$ on X . Assume that G is an algebraic group, and that X is a homogeneous space for G . Assume furthermore that $\mathcal{E}$ is a G -homogeneous vector bundle on X , then it follows that $\mathcal{P}^k(\mathcal{E})$ is again a G -homogeneous vector bundle on X . Consider the line bundle $\mathcal{O}(n)$ on $\mathbf{P}(V^*) = SL(V)/P$, then $\mathcal{O}(n)$ is an $SL(V)$ -homogeneous line bundle on $\mathbf{P}(V^*)$ for all n , and we may consider the $SL(V)$ -homogeneous vector bundle $\mathcal{P}^k(\mathcal{O}(n))$. We want to compute the representation ρ of P corresponding to the homogeneous vector bundle $\mathcal{P}^k(\mathcal{O}(n))$ for all $1 \leq k < n$ on $\mathbf{P}(V^*)$. Let in the following $X = \mathbf{P}(V^*)$ and consider the projection maps p, q from $X \times X$ to X . Let $\mathcal{I}$ in $\mathcal{O}_{X \times X}$ be the ideal of the diagonal. We have an exact sequence of $SL(V)$ -homogeneous vector bundles on $X \times X$:

$$(2.1.1) \quad 0 \rightarrow \mathcal{I}^{k+1} \rightarrow \mathcal{O}_{X \times X} \rightarrow \mathcal{O}_{\Delta^k} \rightarrow 0.$$

Apply the functor $p_*(- \otimes q^* \mathcal{O}(n))$ to the sequence 2.1.1 to get a long exact sequence

$$(2.1.2) \quad 0 \rightarrow p_*(\mathcal{I}^{k+1} \otimes q^* \mathcal{O}(n)) \rightarrow p_* q^* \mathcal{O}(n) \rightarrow \mathcal{P}^k(\mathcal{O}(n)) \\ \rightarrow R^1 p_*(\mathcal{I}^{k+1} \otimes q^* \mathcal{O}(n)) \rightarrow R^1 p_* q^* \mathcal{O}(n) \rightarrow R^1 p_*(\mathcal{O}_{\Delta^k} \otimes q^* \mathcal{O}(n)) \rightarrow \dots$$

of vector bundles. The sequence 2.1.2 is a sequence of vector bundles because all sheaves in the sequence are coherent, and it is a standard fact that a homogeneous coherent sheaf on a homogeneous space is locally free. Since the sequence 2.1.2 is a sequence of vector bundles, we get an exact sequence of P -representations when we pass to the fiber at $\bar{e}$.

Consider the diagram

$$\begin{array}{ccc} \mathrm{Spec}(\kappa(\bar{e})) \times X & \xrightarrow{j} & X \times X \\ \downarrow \pi & & \downarrow p \\ \mathrm{Spec}(\kappa(\bar{e})) & \xrightarrow{i} & X \end{array},$$

then by [4], Chapt. III, Sect.12 we get maps

$$\phi^i : R^i p_*(\mathcal{I}^{k+1} \otimes q^* \mathcal{O}(n))(\bar{e}) \rightarrow R^i \pi_*(j^*(\mathcal{I}^{k+1} \otimes q^* \mathcal{O}(n)))$$

of $\mathcal{O}_X$ -modules. Put for any $\mathcal{O}_{X \times X}$ -module $\mathcal{E}$

$$h^i(y, E) = \dim_{\kappa(y)} H^i(X_y, \mathcal{E}_y),$$

where X_y is the fiber $p^{-1}(y)$ and $\mathcal{E}_y$ is the restriction of $\mathcal{E}$ to X_y . We see that

$$h^i(y, \mathcal{I}^{k+1} \otimes q^* \mathcal{O}(n)) = \dim_{\kappa(y)} H^i(X, \mathfrak{m}_y^{k+1} \otimes \mathcal{O}(n))$$

is a constant function of y for $i = 0, 1, \dots$ for the following reason: Consider the following commutative diagram

$$\begin{array}{ccc} \mathrm{Spec}(\kappa(y)) \times X & \xrightarrow{\tilde{g}} & \mathrm{Spec}(\kappa(gv)) \times X \\ \downarrow j & & \downarrow k \\ X \times X & \xrightarrow{g} & X \times X \end{array}$$

where the action of $SL(V)$ on $X \times X$ is given by $g(x, y) = (gx, gy)$. In general if $G \times Y \xrightarrow{\sigma} Y$ is an algebraic group acting on a scheme Y , and $\mathcal{E}$ is a G -linearized sheaf on Y , then there exists an isomorphism $I : \sigma^* \mathcal{E} \rightarrow p^* \mathcal{E}$ where $p : G \times Y \rightarrow Y$ is the projection map. It follows that for all $g \in G$ we get an isomorphism $g^* \mathcal{E} \cong \mathcal{E}$ of sheaves. Then since $\mathcal{O}(n)$ and $\mathcal{I}^{k+1} \otimes q^* \mathcal{O}(n)$ are $SL(V)$ -homogeneous sheaves, we have an isomorphism

$$\tilde{g}^*(\mathfrak{m}_{gy}^{k+1} \otimes \mathcal{O}(n)) = j^* g^*(\mathcal{I}^{k+1} \otimes q^* \mathcal{O}(n)) = \mathfrak{m}_y^{k+1} \otimes \mathcal{O}(n)$$

hence since $\tilde{g}$ is an isomorphism, we see that we have an isomorphism

$$\mathfrak{m}_y^{k+1} \otimes \mathcal{O}(n) \cong \mathfrak{m}_{gy}^{k+1} \otimes \mathcal{O}(n)$$

of sheaves for all g in $SL(V)$. It follows that

$$\dim_{\kappa(y)} H^i(X, \mathfrak{m}_y^{k+1} \otimes \mathcal{O}(n)) = \dim_{\kappa(gy)} H^i(X, \mathfrak{m}_{gy}^{k+1} \otimes \mathcal{O}(n))$$

for all g in $SL(V)$, hence by [4], chapter III Corr. 12.9, it follows that the maps ϕ^i are isomorphisms for $i = 0, 1, \dots$. Here $\mathfrak{m}_y$ is the sheaf of ideals corresponding to the point y in X . We get an exact sequence

$$(2.1.3) \quad \begin{aligned} 0 \rightarrow H^0(X, \mathcal{O}(n) \otimes \mathfrak{m}_{\bar{e}}^{k+1}) &\rightarrow H^0(X, \mathcal{O}(n)) \rightarrow \mathcal{P}^k(\mathcal{O}(n))(\bar{e}) \\ &\rightarrow H^1(X, \mathcal{O}(n) \otimes \mathfrak{m}_{\bar{e}}^{k+1}) \rightarrow H^1(X, \mathcal{O}(n)) \rightarrow \dots \end{aligned}$$

of P -representations.

Lemma 2.2. *For all $1 \leq k < n$ we have that $H^1(X, \mathcal{O}(n) \otimes \mathfrak{m}_{\bar{e}}^{k+1}) = 0$.*

Proof. Consider the exact sequence 2.1.3. We prove that

$$\dim_F H^0(X, \mathcal{O}(n)) - \dim_F H^0(X, \mathcal{O}(n) \otimes \mathfrak{m}_{\bar{e}}^{k+1}) = \dim_F \mathcal{P}^k(\mathcal{O}(n))(\bar{e}),$$

and then the result follows by counting dimensions. We have that $H^0(X, \mathcal{O}(n))$ equals $S^n(V^*)$, where V^* is the F -vectorspace on the basis $x_0, \dots, x_N$. We also see that $H^0(X, \mathcal{O}(n) \otimes \mathfrak{m}_{\bar{e}}^{k+1})$ equals $m^{k+1}S^{n-(k+1)}(V^*)$ considered as a subvectorspace of $S^n(V^*)$. Here m is the F -vectorspace on the basis $x_1, \dots, x_N$ and $m^{k+1}S^{n-(k+1)}(V^*)$ is the image of the natural map

$$S^{k+1}(m) \otimes S^{n-(k+1)}(V^*) \rightarrow S^n(V^*).$$

Write V^* as the direct sum $Fx_0 \oplus m$. Then it follows that

$$m^{k+1}S^{n-(k+1)}(V^*) = x_0^{n-(k+1)}m^{k+1} \oplus \dots \oplus x_0m^{n-1} \oplus m^n,$$

hence we see that the dimension of $m^{k+1}S^{n-(k+1)}(V^*)$ equals $\sum_{i=k+1}^n \binom{i+N-1}{N-1}$. We also see that the dimension of $S^n(V^*)$ equals $\sum_{i=0}^n \binom{i+N-1}{N-1}$, and it follows that

$$\dim_F S^n(V^*) - \dim_F m^{k+1}S^{n-(k+1)}(V^*) = \sum_{i=0}^k \binom{i+N-1}{N-1}.$$

It follows that

$$\sum_{i=0}^k \binom{i+N-1}{N-1} = \binom{k+N}{N} = \dim_F \mathcal{P}^k(\mathcal{O}(n))(\bar{e}),$$

and we have proved that

$$\dim_F H^0(X, \mathcal{O}(n)) - \dim_F H^0(X, \mathcal{O}(n) \otimes \mathfrak{m}_{\bar{e}}^{k+1}) = \dim_F \mathcal{P}^k(\mathcal{O}(n))(\bar{e}),$$

and the result follows from the fact that the sequence 2.1.3 is exact and that $H^1(X, \mathcal{O}(n)) = 0$ for $n \geq 1$. $\square$

Note that by Lemma 2.2 and the sequence 2.1.3, there exists for all $1 \leq k < n$ an exact sequence of P -representations

$$0 \rightarrow H^0(X, \mathcal{O}(n) \otimes \mathfrak{m}_{\bar{e}}^{k+1}) \rightarrow H^0(X, \mathcal{O}(n)) \rightarrow \mathcal{P}^k(\mathcal{O}(n))(\bar{e}) \rightarrow 0.$$

Since the representation $H^0(X, \mathcal{O}(n) \otimes \mathfrak{m}_{\bar{e}}^{k+1})$ equals $m^{k+1}S^{n-(k+1)}(V^*)$ as subrepresentation of $H^0(X, \mathcal{O}(n)) = S^n(V^*)$, it follows that we have an exact sequence of P -representations

$$(2.2.1) \quad 0 \rightarrow m^{k+1}S^{n-(k+1)}(V^*) \rightarrow S^n(V^*) \rightarrow \mathcal{P}^k(\mathcal{O}(n))(\bar{e}) \rightarrow 0.$$

From the exact sequence

$$0 \rightarrow m \rightarrow V^* \rightarrow V^*/m \rightarrow 0,$$

where m is the F -vectorspace on $x_1, \dots, x_N$, we see that the representation V^*/m is the representation corresponding to the module L^* of P , giving the line bundle $\mathcal{O}(1)$ on $X = SL(V)/P$.

Lemma 2.3. *For all $1 \leq k < n$ there exists a surjective map of P -representations*

$$\phi : S^n(V^*) \rightarrow S^{n-k}(L^*) \otimes S^k(V^*).$$

Proof. Recall that we have chosen a basis $e_0, \dots, e_N$ for V , with the property that $\bar{x}_0$ is a basis for L^* . The P -representation m with basis $x_1, \dots, x_N$ gives an exact sequence

$$0 \rightarrow m \rightarrow V^* \rightarrow L^* \rightarrow 0$$

of P -representations. Define a map

$$\phi : S^n(V^*) \rightarrow S^{n-k}(L^*) \otimes S^k(V^*)$$

as follows: $\phi(f) = \bar{x}_0^{n-k} \otimes \partial_0^{n-k}(f)$, where ∂_0^{n-k} is $n-k$ times partial derivative with respect to the x_0 -variable. Let g be an element of P . Then by induction on the degree of the differential-operator ∂_0^{n-k} and applying the chain-rule for derivation, it follows that

$$\begin{aligned} \phi(gf) &= \bar{x}_0^{n-k} \otimes \partial_0^{n-k}(gf) = \bar{x}_0^{n-k} \otimes a^{-(n-k)} g(\partial_0^{n-k} f) = \\ &= a^{-(n-k)} \bar{x}_0^{n-k} \otimes g(\partial_0^{n-k} f) = g(\bar{x}_0^{-(n-k)} \otimes \partial_0^{n-k} f) = g\phi(f), \end{aligned}$$

and we see that ϕ is P -linear. It is clearly surjective, and the lemma follows. $\square$

Theorem 2.4. *For all $1 \leq k < n$, the representation corresponding to $\mathcal{P}^k(\mathcal{O}(n))$ is $S^{n-k}(L^*) \otimes S^k(V^*)$.*

Proof. By Lemma 2.3 there exists a surjective map of P -representations

$$\phi : S^n(V^*) \rightarrow S^{n-k}(L^*) \otimes S^k(V^*).$$

We claim that $m^{k+1}S^{n-(k+1)}(V^*)$ equals $\ker\phi$: We first prove the inclusion

$$m^{k+1}S^{n-(k+1)}(V^*) \subseteq \ker\phi.$$

Pick a monomial $x_0^{p_0} x_1^{p_1} \dots x_N^{p_N}$ in $m^{k+1}S^{n-(k+1)}(V^*)$, hence $p_0 + \dots + p_N = n$ and $p_0 < n-k$. These monomials form a basis for $m^{k+1}S^{n-(k+1)}(V^*)$. We see that $\partial_0^{n-k}(x_0^{p_0} \dots x_N^{p_N})$ is zero, hence since ϕ is a linear map, it follows that we have an inclusion

$$m^{k+1}S^{n-(k+1)}(V^*) \subseteq \ker\phi$$

of vectorspaces. The reverse inclusion follows from counting dimensions and the fact that ϕ is surjective: We have that

$$\begin{aligned} \dim_F \ker\phi &= \dim_F S^n(V^*) - \dim_F S^{n-k}(L^*) \otimes S^k(V^*) \\ &= \sum_{i=0}^n \binom{i+N-1}{N-1} - \sum_{i=0}^k \binom{i+N-1}{N-1} = \sum_{i=k+1}^n \binom{i+N-1}{N-1}, \end{aligned}$$

and we see that $\dim_F \ker\phi = \dim_F m^{k+1}S^{n-(k+1)}(V^*)$ and it follows that

$$m^{k+1}S^{n-(k+1)}(V^*) = \ker\phi,$$

hence we have an exact sequence of P -representations

$$0 \rightarrow m^{k+1} S^{n-(k+1)}(V^*) \rightarrow S^n(V^*) \rightarrow S^{n-k}(L^*) \otimes S^k(V^*) \rightarrow 0.$$

Using sequence 2.2.1 we get isomorphisms

$$\begin{aligned} \mathcal{P}^k(\mathcal{O}(n))(\bar{e}) &\cong H^0(X, \mathcal{O}(n)) / H^0(X, \mathcal{O}(n) \otimes \mathfrak{m}_{\bar{e}}^{k+1}) \cong \\ S^n(V^*) / m^{k+1} S^{n-(k+1)}(V^*) &\cong S^{n-k}(V^*) \otimes S^k(V^*), \end{aligned}$$

and it follows that $\mathcal{P}^k(\mathcal{O}(n))(\bar{e})$ is isomorphic to $S^{n-k}(L^*) \otimes S^k(V^*)$ as representation. $\square$

Note that the result in Theorem 2.4 is true if $\text{char}(F) > n$.

Corollary 2.5. *For all $1 \leq k < n$, $\mathcal{P}^k(\mathcal{O}(n))$ splits as abstract vector bundle as $\oplus \binom{N+k}{N} \mathcal{O}(n-k)$.*

Proof. Since $S^k(V^*)$ corresponds to the trivial rank $\binom{N+k}{N}$ abstract vector bundle on $\mathbf{P}(V^*)$, and $S^{n-k}(L^*)$ corresponds to the line bundle $\mathcal{O}(n-k)$, the assertion is proved. $\square$

We see that we recover results on the splitting-type of the principal parts obtained in [5], [6], [7] and [8].

Acknowledgements. I would like to thank Michel Brion for an invitation to spend spring 2001 at the Institut Joseph Fourier. Thanks for suggesting the problem, sharing ideas and for answering numerous questions on homogeneous spaces and representation theory. Thanks also to Laurent Manivel for stimulating discussions on the subject. I would also like to thank Mina Teicher for an invitation to the Bar-Ilan University, where parts of this work was done and this note was written. Finally, thanks to Dan Laksov for comments.

REFERENCES

- [1] D. N. Akhiezer, Lie group actions in complex analysis, Aspects of Mathematics, *Vieweg* (1995)
- [2] A. Borel, Linear algebraic groups, Graduate Texts in Mathematics no. 126, *Springer Verlag* (1991)
- [3] J. C. Jantzen, Representations of algebraic groups, Pure and Applied Mathematics no. 131, *Academic Press* (1987)
- [4] R. Hartshorne, Algebraic geometry, Graduate Texts in Mathematics no. 52, *Springer Verlag* (1977)
- [5] H. Maakestad, Modules of principal parts on the projective line, *preprint math.AG/0111149*
- [6] D. Perkinson, Principal parts of linebundles on toric varieties, *Compositio Math.* 104, 27-39, (1996)
- [7] R. Piene, G. Sacchiero, Duality for rational normal scrolls, *Comm. in Alg.* 12 (9), 1041-1066, (1984)
- [8] S. di Rocco, A. J. Sommese, Line bundles for which a projectivized jet bundle is a product, *Proc. Amer. Math. Soc.* 129, (2001), no.6, 1659-1663

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, RAMAT GAN, ISRAEL
E-mail address: makesth@macs.biu.ac.il